

M00) a)  $\log_2 5 = \frac{\log_{10} 5}{\log_{10} 2} = \boxed{\frac{1}{x}}$

b)  $\log_2 z = \log_2(z^2 \cdot 5) = 2 \log_2 z + \log_2 5 = \boxed{2x + y}$

M00)  $(a+b)^{12} = \sum_{r=0}^{12} \binom{12}{r} a^r b^{12-r}$

$$\binom{12}{5} a^5 b^7 = \frac{12!}{5! 7!} a^5 b^7 = \boxed{792 a^5 b^7}$$

M00)  $4x^2 + 4kx + 9 = 0 ; (k > 0)$   
 $x = \frac{-4k \pm \sqrt{16k^2 - 144}}{8}$

Solucionando  $\Rightarrow 16k^2 = 144 ; k^2 = \frac{144}{16} ; k = \pm \sqrt{\frac{144}{16}}$   $\begin{cases} \frac{12}{4} = 3 \\ -\frac{12}{4} = -3 \end{cases}$  Dice que  $k > 0$

N00) a)  $C = 1000 \cdot (1 + 0.075)^{10} = 1000 \cdot 1.075^{10} = \boxed{2061,03 \$}$

b)  $C = 1000 \cdot (1.075^{10} + 1000 \cdot 1.075^9 + \dots + 1000 \cdot 1.075^1) = \begin{matrix} \text{(Suma de los términos} \\ \text{de una P.G. con r = 1.075)} \end{matrix}$   
 $= \frac{1000 \cdot 1.075^{10} \cdot 1.075 - 1000 \cdot 1.075}{1.075 - 1} = \frac{1000 \cdot 1.075 \cdot (1.075^{10} - 1)}{0.075} = \boxed{15208,12 \$}$

N00)  $V = 10000 \cdot 0.933^t$

a)  $t=0 \Rightarrow V = 10000 \text{ l}$

b)  $V = 5000 \Rightarrow 5000 = 10000 \cdot 0.933^t ; 0.5 = 0.933^t$

$$t = \log_{0.933} 0.5 = \frac{\log 0.5}{\log 0.933} = 9.9949 \text{ min} \approx 9 \text{ min } 60 \text{ seg} \approx \boxed{10 \text{ min}}$$

c) 5% de 10.000 = 500

$$500 = 10000 \cdot 0.933^t ; 0.05 = 0.933^t$$

$$t = \log_{0.933} 0.05 = \frac{\log 0.05}{\log 0.933} = 43,20 \text{ min} \approx \boxed{45 \text{ min}} \checkmark$$

M01) a)  $P = 1512 \cdot (1 + 0.027)^{-1} = \frac{1512}{1.027} = \boxed{148 \text{ millions}}$

b)  $P = 1512 \cdot 1.027^5 = \frac{1512}{1.027^5} = \boxed{133 \text{ millions}}$

M01) a)  $C = 1500 \cdot (1 + 0.0525)^3 = 1748,87 \approx \boxed{1749 \text{ f.}}$

b)  $3000 = 1500 \cdot 1.00525^t ; 2 = 1.00525^t$

$$t = \log_{1.00525} 2 = \frac{\log 2}{\log 1.00525} = 132,37 \rightarrow \boxed{133 \text{ años}}$$

M01)  $3000 = 1500 \cdot (1+i)^{10} ; 2 = (1+i)^{10} ; 1+i = \sqrt[10]{2} ; i = \sqrt[10]{2} - 1 = 0.0718 = \boxed{7,18 \%}$

$$N_01 \quad C = 5000 \cdot (1+0.065)^5 = 6850,43 \approx \boxed{6850,43}$$

$$N_01 \quad a) \quad N = 5000 \cdot e^{kt}$$

$$\frac{N=2500}{t=s} \Rightarrow 2500 = 5000 \cdot e^{-sk} ; \quad 0.5 = e^{-sk} ; \quad -sk = \ln 0.5 ; \quad k = \frac{\ln 0.5}{-s} = \boxed{0.139}$$

$$b) \quad N = 50 : \quad 50 = 5000 \cdot e^{-0.139t} ; \quad 0.01 = e^{-0.139t} ;$$

$$-0.139t = \ln 0.01 ; \quad t = \frac{\ln 0.01}{-0.139} = 33.13 \dots \quad \boxed{33 \text{ años}}$$

$$N_01 \quad (x^3 - 3y^2)^5 = \sum_{r=0}^5 \binom{5}{r} (x^3)^r (-3y^2)^{5-r} = \sum_{r=0}^5 \binom{5}{r} x^{3r} (-3)^{5-r} y^{10-2r}$$

$$\text{Exponents iguales} \Rightarrow 3r = 10 - 2r ; \quad 5r = 10 ; \quad r = 2$$

$$r=2 \Rightarrow \binom{5}{2} \cdot (x^3)^2 \cdot (-3y^2)^{5-2} = 10 \cdot x^6 \cdot (-3)^3 \cdot y^6 = \boxed{-270x^6y^6}$$

$$N_01 \quad \log_9 81 + \log_9 \left(\frac{1}{9}\right) + \log_9 3 = \log_9 x$$

$$81 \cdot \frac{1}{9} \cdot 3 = x ; \quad \boxed{x = 27}$$

$$M_02 \quad a) \quad \left(3x^2 - \frac{1}{x}\right)^9 \text{ tiene } \boxed{dici3} \text{ sumandos.}$$

$$b) \quad \left(3x^2 - \frac{1}{x}\right)^9 = \sum_{r=0}^9 \binom{9}{r} \cdot (3x^2)^r \left(-\frac{1}{x}\right)^{9-r} = \sum_{r=0}^9 \binom{9}{r} \frac{3^r x^{2r} \cdot (-1)^{9-r}}{x^{9-r}}$$

$$2r = 9 - r \Rightarrow r = 3$$

$$r=3 \rightarrow \binom{9}{3} (3x^2)^3 \left(-\frac{1}{x}\right)^6 = 84 \cdot 27 \cdot x^6 \cdot \frac{1}{x^6} = \boxed{2268}$$

$$M_02 \quad \log_{27} x = 1 - \log_{27} (x-0.4) ; \quad \log_{27} x + \log_{27} (x-0.4) = 1 ; \quad \log_{27} x(x-0.4) = 1$$

$$x(x-0.4) = 27^1 ; \quad x^2 - 0.4x = 27 ; \quad x^2 - 0.4x - 27 = 0$$

$$x = \frac{0.4 \pm \sqrt{0.16 + 108}}{2} = \frac{0.4 \pm 10.4}{2} \quad \boxed{s^4}$$

No existen soluciones para los logaritmos

$$(2-x)^5 = \sum_{r=0}^5 \binom{5}{r} 2^r (-x)^{5-r}$$

$$5-r=3 \Rightarrow r=2 \rightarrow \binom{5}{2} \cdot 2^2 \cdot (-x)^3 = 10 \cdot 4 \cdot (-x^3) = \boxed{-40x^3}$$

$$N_02 \quad 3000 = 10000 \cdot \left(1 + \frac{0.15}{12}\right)^t ; \quad 3 = \left(1 + 0.0125\right)^t ; \quad 3 = 1.0125^t ;$$

$$t = \log_{1.0125} 3 = \frac{\log 3}{\log 1.0125} = 88,44 \rightarrow \boxed{89 \text{ meses}}$$

$$M_03 \quad (5+2x^2)^7 = \sum_{r=0}^7 \binom{7}{r} 5^r (2x^2)^{7-r} = \sum_{r=0}^7 \binom{7}{r} 5^r 2^{7-r} x^{14-2r}$$

$$14-2r=10 ; \quad r=2 \rightarrow \binom{7}{2} 5^2 (2x^2)^5 = 21 \cdot 25 \cdot 32 \cdot x^{10} = \boxed{16800x^{10}}$$

$$M03 \quad a) \log_5 x^2 = 2 \log_5 x = \boxed{2y}$$

$$b) \log_5 \left(\frac{1}{x}\right) = \log_5 1 - \log_5 x = -\log_5 x = \boxed{-y}$$

$$c) \log_{25} x = \frac{\log_5 x}{\log_5 25} = \frac{\log_5 x}{\log_5 5^2} = \boxed{\frac{y}{2}}$$

$$N03 \quad Kx^2 + 3x + 1 = 0 \\ x = \frac{-3 \pm \sqrt{9 - 4K}}{2K}$$

$$\text{Unique solution} \Rightarrow 9 - 4K = 0 ; \boxed{K = 9/4}$$

$$N03 \quad m = 4 e^{-0.2t}$$

$$a) t=0 \Rightarrow \boxed{m=4 \text{ kg}}$$

$$b) 1's = 4 e^{-0.2t} ; 0'375 = e^{-0.2t} ; -0.2t = \ln 0'375 ;$$

$$t = \frac{\ln 0'375}{-0.2} = 8'90 \text{ hours} = \boxed{8 \text{ hours } 54 \text{ min}}$$

$$N03 \quad (2+ax)^4 = 2^4 + 4 \cdot 2^3 \cdot ax + 6 \cdot 2^2 \cdot (ax)^2 + 4 \cdot 2 \cdot (ax)^3 + (ax)^4 = \\ = \boxed{16 + 32ax + 24a^2x^2 + 8a^3x^3 + a^4x^4}$$

$$M04 \quad b) (1+x^2)^6 = \sum_{r=0}^6 \binom{6}{r} 1^r (x^2)^{6-r} = \\ = \sum_{r=0}^6 \binom{6}{r} x^{12-2r}$$

$$12-2r=8 \Rightarrow \boxed{r=2}$$

$$\binom{6}{2} x^8 = \boxed{15x^8}$$

$$a) \begin{array}{ccccccc} & & 1 & 1 & & & \\ & & 1 & 2 & 1 & & \\ & & 1 & 3 & 3 & 1 & \\ & & 1 & 4 & 6 & 4 & 1 \\ & & 1 & 5 & 10 & 10 & 5 & 1 \\ & & 1 & 6 & 15 & 20 & 15 & 6 & 1 \end{array}$$

$$M04 \quad a) e^{\ln x} = \boxed{x} \quad b) e^{\ln x + \ln y} = e^{\ln x} \cdot e^{\ln y} = \boxed{xy}$$

$$c) \ln(e^{x+y})^2 = 2 \ln e^{x+y} = 2(x+y) = \boxed{2x+2y}$$

$$M04 \quad \log_{10} \left( \frac{x}{y^2 \sqrt{z}} \right) = \log_{10} x + 2 \log_{10} y + \frac{1}{2} \log_{10} z = \boxed{p - 2q - \frac{r}{2}}$$

$$M04 \quad x^2 - 2kx + 1 = 0 \\ x = \frac{2k \pm \sqrt{4k^2 - 4}}{2}$$

Dois raios reais distintos  $\Rightarrow 4k^2 - 4 > 0 ; 4k^2 > 4 ; k^2 > 1$

$$k^2 = 1 \rightarrow \boxed{k = \pm 1}$$

$$\begin{array}{ccccc} & -1 & & +1 & \\ 4k^2 - 4 & + & - & + & \end{array}$$

$$\boxed{k \in (-\infty, -1) \cup (1, +\infty)}$$

M04

$$(2+ax)^{10} = \sum_{r=0}^{10} \binom{10}{r} 2^r (ax)^{10-r}$$

$$10-r=3 \rightarrow r=7 \quad \binom{10}{7} 2^7 (ax)^3 = 120 \cdot 128 \cdot a^3 x^3 = \frac{15360 a^3 x^3}{\approx 414720}$$

$$15360 a^3 = 414720 \Rightarrow a^3 = 27 \Rightarrow a = 3$$

M04

(i) Comienzos de 2004  $\rightarrow n=10 \rightarrow P = 1200000 \cdot 1.025^{10} \approx 1536101 \text{ hab}$

(ii)  $\frac{1536101}{1200000} = 1.28 = 128\% \rightarrow \text{Creció un } 28\%$

(iii)  $P > 2000000 \Rightarrow 2000000 = 1.200000 \cdot 1.025^m$

$$\frac{2000000}{1200000} = 1.025^m$$

$$1.6 = 1.025^m ; m = \log_{1.025} 1.6 = 20.687 \Rightarrow m=21 \rightarrow 2015$$

N04

$$(2-3x)^8 = \sum_{r=0}^8 \binom{8}{r} 2^r (-3x)^{8-r}$$

$$8-r=3 \rightarrow r=5 \quad \binom{8}{5} \cdot 2^5 \cdot (-3x)^3 = 56 \cdot 32 \cdot (-27)x^3 = 48384x^3$$

N04

$$\log \frac{x^2 \sqrt{y}}{z^3} = 2 \log x + \frac{1}{2} \log y - 3 \log z = 2a + \frac{b}{2} - 3c$$

N04

a) Sueldo 2º año = 11000 + 400 = 11400 \$  
 " 3º año = 11400 + 400 = 11800 \$

$$\begin{aligned} \text{Total} &= 11000 + 11400 + 11800 + \dots + (11000 + 4 \cdot 400) = \\ &= 11000 + 11400 + \dots + 14600 = \text{(Suma 10 términos)} \\ &= \frac{11000 + 14600}{2} \cdot 10 = 128000 \$ \end{aligned}$$

b) Sueldo 2º año = 10000 + 10000 \cdot 1.07 = 10000 \cdot 1.07 = 10700 \\$  
 Sueldo 3º año = 10700 + 10700 \cdot 1.07 = 10700 \cdot 1.07 = 11449 \\$  
 Sueldo 10º año = 10000 \cdot 1.07^9 = 18384,59 \\$

c) Sueldos en n años iniciales:  $S_A = 10000 + (n-1)400$

$$S_B = 10000 \cdot 1.07^{n-1}$$

Totales en n años:  $\text{Total A} = \frac{10000 + (10000 + (n-1)400)}{2} \cdot n = 10800n + 200n^2$

$$\text{Total B} = \frac{10000 \cdot 1.07^{n-1} - 1.07 - 10000}{1.07 - 1} = 142857,14 \cdot (1.07^n - 1)$$

$$\text{Total B} > \text{Total A} \Rightarrow 142857,14 (1.07^n - 1) > 10800n + 200n^2$$

Resuelto con calculadora gráfica  $\rightarrow n = 6,33 \Rightarrow n = 7 \text{ años}$

Por tanto:  $m=3$ : Total A = 34200 \\$  
 Total B = 32149 \\$  
 $m=4$ : Total A = 46400 \\$  
 Total B = 44399,43 \\$

$$\left\{ \begin{array}{l} m=5: \text{Total A} = 59000 \$ \\ \text{Total B} = 57507,39 \$ \\ m=6: \text{Total A} = 72000 \$ \\ \text{Total B} = 71532,91 \$ \\ m=7: \text{Total A} = 85400 \$ \\ \text{Total B} = 86540,21 \$ \end{array} \right.$$

$$M05 \quad (x^2 - 2)^5$$

a) Son seis términos

b)  $(x^2 - 2)^5 = \sum_{r=0}^5 \binom{5}{r} (x^2)^r (-2)^{5-r}$

$$r=2 \rightarrow \binom{5}{2} (x^2)^2 (-2)^3 = 10x^4 \cdot (-8) = -80x^4 \rightarrow A = -80$$

$$M05 \quad 9^{2x} = 27^{1-x}; \quad (3^2)^{2x} = (3^3)^{1-x}; \quad 3^{4x} = 3^{3-x}; \quad 4x = 3-x; \quad 5x = 3; \quad x = 3/5$$

$$M05 \quad a) \log_3 x - \log_3 (x-5) = \log_3 \frac{x}{x-5} \Rightarrow A = \frac{x}{x-5}$$

$$b) \log_3 x - \log_3 (x-5) = 1; \quad \log_3 \frac{x}{x-5} = 1; \quad \frac{x}{x-5} = 3^1; \quad x = 3x - 15; \quad -2x = -15 \quad |x = 15/2$$

$$N05 \quad (3 + \sqrt{7})^3 = 3^3 + 3 \cdot 3^2 \sqrt{7} + 3 \cdot 3 (\sqrt{7})^2 + (\sqrt{7})^3 = 27 + 27\sqrt{7} + 63 + 7\sqrt{7} = \\ = 90 + 34\sqrt{7} \rightarrow p = 90 \quad q = 24$$

$$N05 \quad V = 10000 e^{-0.3t}$$

$$V < 1500 \rightarrow 1500 = 10000 e^{-0.3t}; \quad 0.15 = e^{-0.3t}; \quad -0.3t = \ln 0.15; \\ t = \frac{\ln 0.15}{-0.3} = 6.32 \rightarrow t = 7 \text{ años}$$

Muestre  
06/08

a)  $C = 5000 \cdot 1.063^m$

b)  $C_5 = 5000 \cdot 1.063^5 = 6786,35 \$$

c)  $10000 < 5000 \cdot 1.063^m$

$$2 < 1.063^m; \quad m > \log_{1.063} 2 = 11.35 \rightarrow m = 12 \text{ años}$$

$$M06 \quad a) \log_c 15 = \log_c (3 \cdot 5) = \log_c 3 + \log_c 5 = p+q$$

$$\log_c 25 = \log_c 5^2 = 2 \log_c 5 = 2q$$

$$b) \log_d 6 = \frac{1}{2} \rightarrow 6 = d^{1/2}; \quad 6 = \sqrt{d}; \quad d = 6^2; \quad d = 36$$

$$M06 \quad a) \ln(x+2) = 3; \quad x+2 = e^3; \quad x = e^3 - 2$$

$$b) 10^{2x} = 500; \quad 2x = \log 500; \quad x = \frac{\log 500}{2} = 1.35$$

$$N06 \quad a) 5^{x+1} = 625; \quad x+1 = 4; \quad x = 3$$

$$b) \log_a (3x+5) = 2; \quad 3x+5 = a^2; \quad 3x = a^2 - 5; \quad x = \frac{a^2 - 5}{3}$$

$$N06 \quad a) \ln a^3 b = 3 \ln a + \ln b = 3p+q$$

$$b) \ln \frac{\sqrt{a}}{b} = \frac{1}{2} \ln a - \ln b = \frac{p}{2} - q$$

M07

$$\underline{a} \quad N = 250000 \cdot 10^{13} = \boxed{253250 \text{ hab}}$$

$$\underline{b} \quad 2002 - 1972 = 30$$

$$N = 250000 \cdot 10^{13}^{30} = \boxed{368318 \text{ hab}}$$

M07

$$(x+2y)^{10} = \sum_{r=0}^{10} \binom{10}{r} x^r (2y)^{10-r}$$

$$\binom{10}{8} x^8 (2y)^2 = 45x^8 \cdot 4y^2 = 180x^8 y^2 ; \quad \boxed{a=180}$$

M07

$$\underline{a} \quad \log_{10} 10 = \log_{10}(5 \cdot 2) = \log_{10} 5 + \log_{10} 2 = \boxed{1+9}$$

$$\underline{b} \quad \log_8 8 = \log_8 2^3 = 3 \log_8 2 = \boxed{39}$$

$$\underline{c} \quad \log_{10} 25 = \log_{10} \left(\frac{5}{2}\right) = \log_{10} 5 - \log_{10} 2 = \boxed{1-9}$$

M07

$$(x^3 - 3x)^6 = \sum_{r=0}^6 \binom{6}{r} (x^3)^r (-3x)^{6-r} = \sum_{r=0}^6 \binom{6}{r} (-3)^{6-r} x^{3r+6-r} =$$

$$= \sum_{r=0}^6 \binom{6}{r} (-3)^{6-r} x^{2r+6}$$

a Time + terminos

$$\underline{b} \quad 2r+6=12 ; \quad 2r=6 ; \quad \underbrace{r=3} \rightarrow \binom{6}{3} (-3)^3 x^{12} = 20 \cdot (-27) x^{12} = \boxed{-540x^{12}}$$

M07

$$\underline{a} \quad (2^x+a)(2^x+b) = (2^x)^2 + a \cdot 2^x + b \cdot 2^x + ab = (2^x)^2 + \underbrace{(a+b)}_{ab=-12} 2^x + \underbrace{ab}_{= (2^x)^2 + 2^x - 12} =$$

$$\begin{array}{l} a+b=1 \xrightarrow{b=1-a} \\ ab=-12 \end{array} \rightarrow a(1-a)=-12 ; \quad a-a^2=-12 ; \quad 0=a^2-a-12 ;$$

$$\left[ a = \frac{1 \pm \sqrt{1+48}}{2} = \frac{1 \pm 7}{2} \right] \quad \left\{ \begin{array}{l} 4 \rightarrow b=-3 \\ -3 \rightarrow b=4 \end{array} \right\} \Rightarrow (2^x)^2 + 2^x - 12 = (2^x+4)(2^x-3)$$

$$\underline{b} \quad (2^x)^2 + 2^x - 12 = 0 ; \quad (2^x+4)(2^x-3) = 0 \quad \begin{array}{l} 2^x+4=0 ; \quad 2^x=-4 ; \quad \text{sin solucion.} \\ 2^x-3=0 ; \quad 2^x=3 ; \quad \boxed{x = \log_2 3} \end{array}$$

Sólo tiene una solución porque las exponentiales tienen valores mayores que 0.

M07

$$\underline{a} \quad \left(e + \frac{1}{e}\right)^4 = e^4 + 4e^3 \frac{1}{e} + 6e^2 \left(\frac{1}{e}\right)^2 + 4e \left(\frac{1}{e}\right)^3 + \left(\frac{1}{e}\right)^4 =$$

$$= \boxed{e^4 + 4e^2 + 6 + 4e^{-2} + e^{-4}}$$

$$\underline{b} \quad \left(e + \frac{1}{e}\right)^4 + \left(e - \frac{1}{e}\right)^4 = (e^4 + 4e^2 + 6 + 4e^{-2} + e^{-4}) + (e^4 - 4e^2 + 6 - 4e^{-2} + e^{-4}) =$$

$$= \boxed{2e^4 + 12 + 2e^{-4}}$$

Muestra 08

a)  $\log_x 49 = 2$ ;  $x^2 = 49$ ;  $x = \sqrt{49}$  la base no puede ser negativa

b)  $\log_2 8 = x$ ;  $x = \log_2 2^3 = 3$

c)  $\log_{25} x = -\frac{1}{2}$ ;  $25^{-\frac{1}{2}} = x$ ;  $x = \frac{1}{\sqrt{25}} = \frac{1}{5}$

d)  $\log_2 x + \log_2 (x-7) = 3$

$\log_2 x(x-7) = 3$ ;  $x(x-7) = 2^3$ ;  $x^2 - 7x - 8 = 0$ ;  $x^2 - 7x - 8 = 0$

$$x = \frac{7 \pm \sqrt{49 + 32}}{2} = \frac{7 \pm 9}{2}$$

M08

$\left(\frac{2}{3}x - 3\right)^8 = \sum_{r=0}^8 \binom{8}{r} \left(\frac{2}{3}x\right)^r \cdot (-3)^{8-r}$

(r=3)  $\rightarrow \binom{8}{3} \left(\frac{2}{3}x\right)^3 (-3)^5 = 56 \cdot \frac{8}{27} x^3 \cdot (-243) = -4032 x^3$

M08

T = 280 · 1/12<sup>m</sup>

a) m=5  $\rightarrow T = 280 \cdot 1/12^5 = 493,4 \approx 493$  personas

T = 560  $\rightarrow 560 = 280 \cdot 1/12^m$ ;  $2 = 1/12^m$ ;  $m = \log_2 12 \approx 4,1$   $\rightarrow 2007$

b) m=5  $\rightarrow P = \frac{2560000}{10 + 90 e^{-0,1m}} \approx 139636$  personas

$$2 \cdot 256000 = \frac{2560000}{10 + 90 e^{-0,1m}}; 10 + 90 e^{-0,1m} = \frac{2560000}{2 \cdot 25600}; 10 + 90 e^{-0,1m} = 50;$$

$$90 e^{-0,1m} = 40; e^{-0,1m} = \frac{4}{9}; -0,1m = \ln \frac{4}{9}; m = \frac{\ln \frac{4}{9}}{-0,1} = 8,109 \Rightarrow \underline{\underline{m}}$$

c) m=0: R =  $\frac{25600}{280} = 91,43$

$$R < 70 \rightarrow 70 > \frac{P}{T} = \frac{2560000}{280 \cdot 1/12^m}; 19600 \cdot 1/12^m > \frac{2560000}{10 + 90 e^{-0,1m}};$$

$$196 \cdot 1/12^m > \frac{2560}{1 + 9e^{-0,1m}}$$

con calculadora gráfica:  $m = 9,31 \rightarrow 10$  años

N08

a)  $(x-2)^4 = x^4 + 4x^3(-2) + 6x^2(-2)^2 + 4x(-2)^3 + (-2)^4 =$   
 $= x^4 - 8x^3 + 24x^2 - 32x + 16$

b)  $(3x+4) \cdot (x-2)^4 = (3x+4) \underbrace{(x^4 - 8x^3 + 24x^2 - 32x + 16)}_{\uparrow \uparrow}$

$$3x^4 + 4 \cdot (-8) = 140$$

a) m=10

b) a=p; b=q

c)  $\binom{10}{5} p^5 q^5$

M09

M09

$$\text{a)} \log_2 32 = \log_2 2^5 = \boxed{5}$$

$$\text{b)} \log_2(32^x / 8^y) = x \log_2 32 - y \log_2 8 = 5x - 3y \rightarrow \boxed{x=5} ; \boxed{y=-3}$$

M09

$$\text{c)} kx^2 + (k-3)x + 1 = 0$$

$$x = \frac{-(k-3) \pm \sqrt{(k-3)^2 - 4k}}{2k}$$

Solutions iguales  $\Rightarrow (k-3)^2 - 4k = 0 ; k^2 - 6k + 9 - 4k = 0 ; k^2 - 10k + 9 = 0 ;$

$$k = \frac{10 \pm \sqrt{100-36}}{2} = \frac{10 \pm 8}{2} \begin{array}{|c|} \hline 9 \\ \hline 1 \\ \hline \end{array}$$

$$\text{b)} x^2 + (k-3)x + k = 0$$

$$x = \frac{-(k-3) \pm \sqrt{(k-3)^2 - 4k}}{2k}$$

Solutions iguales  $\Rightarrow (k-3)^2 - 4k = 0 \rightarrow \boxed{k=1}$

M10

$$\text{a)} (2+x)^7 = 2^7 + 7 \cdot 2^6 x + 21 \cdot 2^5 x^2 + 35 \cdot 2^4 x^3 + \dots = \boxed{16 + 32x + 24x^2 + 8x^3 + x^4}$$

$$\text{b)} (2+x)^7 \cdot \left(1 + \frac{1}{x^2}\right) = \left(16 + 32x + 24x^2 + 8x^3 + x^4\right) \left(1 + \frac{1}{x^2}\right)$$

$$24+1 = \boxed{25}$$

M10

$$\log_2 x + \log_2(x-3) = 3 ; \log_2 x(x-3) = 3 ; x(x-3) = 2^3 ; x^2 - 3x = 8 ; x^2 - 3x - 8 = 0$$

$$x = \frac{3 \pm \sqrt{9+32}}{2} = \begin{array}{|c|} \hline \frac{3+\sqrt{41}}{2} \\ \hline \cancel{\frac{3-\sqrt{41}}{2}} \\ \hline \end{array} \quad \text{Dado que } x > 2$$

M10

$$(3x^2 - \frac{2}{x})^5 = \sum_{r=0}^5 \binom{5}{r} (3x^2)^r \left(-\frac{2}{x}\right)^{5-r} = \sum_{r=0}^5 \binom{5}{r} 3^r (-2)^{5-r} \cdot \frac{x^{2r}}{x^{5-r}} = \sum_{r=0}^5 \binom{5}{r} 3^r (-2)^{5-r} \cdot x^{3r-5}$$

$$3r-5=4 ; 3r=9 ; r=3 \rightarrow \binom{5}{3} (3x^2)^3 \left(\frac{-2}{x}\right)^2 = 10 \cdot 27 \cdot x^6 \cdot \frac{4}{x^2} = \boxed{1080x^4}$$

M11

$$(x+2)^{11} = \sum_{r=0}^{11} \binom{11}{r} x^r 2^{11-r}$$

a) Son doce términos.

$$\text{b)} \quad r=2 \rightarrow \binom{11}{2} x^2 2^9 = \boxed{28160x^2}$$

N11

$$\text{a)} A(0) = \boxed{10 \text{ mg/l}}$$

$$\text{b)} A(50) = 10 \cdot 0.5^{0.014 \cdot 50} = \boxed{6.16 \text{ mg/l}}$$

$$\text{c)} 0.395 = 10 \cdot 0.5^{0.014t} ; 0.395 = 0.5^{0.014t} ; 0.014t = \log_{0.5} 0.395 ;$$

$$t = \frac{\log 0.395}{0.014} = 333 \text{ min} = 5 \text{ h } 33 \text{ min} \rightarrow \boxed{18:33}$$

$$N11 \quad (3x^2+2)^9 = \sum_{r=0}^9 \binom{9}{r} (3x^2)^r (2)^{9-r}$$

a) Hay diez términos

b)  $\binom{9}{2} \rightarrow \binom{9}{2} (3x^2)^2 2^7 = 36 \cdot 9x^4 \cdot 128 = \boxed{41472x^4}$

$$M12 \quad x^2 + (k-1)x + 1 = 0$$

$$x = \frac{-(k-1) \pm \sqrt{(k-1)^2 - 4}}{2}$$

Raíces iguales  $\Rightarrow (k-1)^2 - 4 = 0 ; k^2 - 2k + 1 - 4 = 0 ; k^2 - 2k - 3 = 0 ;$

$$k = \frac{2 \pm \sqrt{4+4}}{2} = \frac{2 \pm 4}{2} \begin{cases} 3 \\ -1 \end{cases}$$

$$N12 \quad \left(2x^3 + \frac{b}{x}\right)^8 = \sum_{r=0}^8 \binom{8}{r} (2x^3)^r \left(\frac{b}{x}\right)^{8-r} = \sum_{r=0}^8 \binom{8}{r} 2^r \cdot b^{8-r} x^{4r-8}$$

a)  $\binom{8}{7} \rightarrow \binom{8}{7} (2x^3)^7 \cdot \frac{b}{x} = 8 \cdot 128 \cdot x^21 \cdot \frac{b}{x} = \boxed{1024b x^{21}} = 3072$

$$1024b = 3072 \rightarrow \boxed{b = 3}$$

b)  $\binom{8}{2} \rightarrow \binom{8}{2} (2x^3)^2 \left(\frac{3}{x}\right)^6 = 28 \cdot 4x^6 \cdot \frac{729}{x^6} = 81648 ; \boxed{k = 81648}$

$$N12 \quad x^2 - 3x + k^2 = 0 ; \quad x^2 - 3x + (k^2 - 4) = 0$$

$$x = \frac{3 \pm \sqrt{9 - 4(k^2 - 4)}}{2}$$

Dois raízes reais distintas  $\Rightarrow 9 - 4(k^2 - 4) > 0 ; 25 - 4k^2 > 0$

$$25 - 4k^2 = 0 ; 25 = 4k^2 ; k = \pm \frac{5}{2} \quad \boxed{25 - 4k^2 = \begin{array}{|c|c|} \hline -\frac{5}{2} & \frac{5}{2} \\ \hline \end{array}} \quad \boxed{k \in (-\frac{5}{2}, \frac{5}{2})}$$

$$M12 \quad \left(1 + \frac{2}{3}x\right)^m (3+mx)^2 = 9 + 84x + \dots$$

$$\left[1 + \binom{m}{1} 1^{m-1} \left(\frac{2}{3}x\right) + \dots\right] \cdot [9 + 6mx + \dots] = 9 + 84x + \dots$$

$$\left[1 + \frac{2m}{3}x + \dots\right] \cdot [9 + 6mx + \dots] = 9 + 84x + \dots$$

$$\left[9 + 6mx + \frac{18mx}{3} + \dots\right] = 9 + 84x + \dots$$

$$\left[9 + \underbrace{12mx}_{12m} + \dots\right] = 9 + 84x + \dots ; 12m = 84 ; \boxed{m = 7}$$

$$N12 \quad (2x+p)^6 = \sum_{r=0}^6 \binom{6}{r} (2x)^r p^{6-r}$$

$\binom{6}{4} \rightarrow \binom{6}{4} (2x)^4 p^2 = 15 \cdot 16x^4 p^2 = \underbrace{240p^2 x^4}_{=60} ; 240p^2 = 60 ; p^2 = \frac{1}{4} ; \boxed{p = \pm \frac{1}{2}}$

$$M13 \quad a) \log_3 p^2 = 2 \log_3 p = 2 \cdot 6 = \boxed{12}$$

b)  $\log_3 \left(\frac{p}{q}\right) = \log_3 p - \log_3 q = 6 - 7 = \boxed{-1}$

c)  $\log_3 (qp) = \log_3 q + \log_3 p = 2 + 6 = \boxed{8}$

$$\textcircled{M13} \quad \triangleq \log_2 40 - \log_2 5 = \log_2 \frac{40}{5} = \log_2 8 = \boxed{3}$$

$$\underline{\text{b)} \quad 8^{\log_2 5} = (2^3)^{\log_2 5} = 2^{3 \log_2 5} = 2^{\log_2 5^3} = 5^3 = \boxed{125}$$

$$\textcircled{M13} \quad \left(\frac{x}{a} + \frac{a^2}{x}\right)^6 = \sum_{r=0}^6 \binom{6}{r} \left(\frac{x}{a}\right)^r \cdot \left(\frac{a^2}{x}\right)^{6-r} = \sum_{r=0}^6 \binom{6}{r} \frac{1}{a^r} \cdot a^{12-2r} \cdot x^{2r-6}$$

$$2r-6=0 ; \quad r=\underline{\underline{3}} \quad \rightarrow \quad \binom{6}{3} \left(\frac{x}{a}\right)^3 \left(\frac{a^2}{x}\right)^3 = 20 \frac{x^3}{a^3} \cdot \frac{a^6}{x^3} = \underline{\underline{20a^3}}$$

$$20a^3 = 1280 ; \quad a^3 = 64 ; \quad a = \sqrt[3]{64} = \boxed{4}$$

$$\textcircled{M13} \quad (3x-2)^{12} = \sum_{r=0}^{12} \binom{12}{r} (3x)^r \cdot (-2)^{12-r}$$

$$\text{a)} \quad \boxed{r=5 \rightarrow p=5 ; q=7}$$

$$\underline{\text{b)} \quad \binom{12}{5} (3x)^5 (-2)^7 = 792 \cdot 243 x^5 \cdot (-128) = \boxed{-24634368 x^5}$$

M 14

a) ii)  $\log_3 27 = \log_3 3^3 = \boxed{3}$

iii)  $\log_8 \frac{1}{8} = \log_8 8^{-1} = \boxed{-1}$

iv)  $\log_{16} \sqrt{16} = \log_{16} 16^{1/2} = \boxed{\frac{1}{2}}$

b)  $3 - 1 - \frac{1}{2} = \log_n x \Rightarrow \frac{3}{2} = \log_n x \Rightarrow \boxed{4^{\frac{3}{2}} = x} \Rightarrow x = \sqrt[4]{4^3}$

M 14

a)  $\log_6 36 = \log_6 6^2 = \boxed{2}$

b)  $\log_6 6 + \log_6 9 = \log_6 36 = \boxed{2}$

c)  $\log_6 2 - \log_6 12 = \log_6 \frac{2}{12} = \log_6 \frac{1}{6} = \log_6 6^{-1} = \boxed{-1}$

M 14

a)  $A(0) = 12 \cdot e^{0 \cdot h \cdot 0} = \boxed{12}$  bacteria

b)  $A(h) = 12 \cdot e^{0 \cdot h \cdot h} = 12 \cdot e^{h^2} \approx 59^{h^2} \approx \boxed{59}$  bacteria

c)  $400 = 12 \cdot e^{0 \cdot h \cdot t} \Rightarrow \ln\left(\frac{400}{12}\right) = 0 \cdot h \cdot t \cdot \ln e \Rightarrow t = \frac{\ln\left(\frac{400}{12}\right)}{0 \cdot h} \approx \boxed{877}$  hours

d)  $60 = 24 \cdot e^{K \cdot 4} \Rightarrow \ln\left(\frac{60}{24}\right) = 4K \cdot \ln e \Rightarrow K = \frac{\ln\left(\frac{60}{24}\right)}{4} \approx \boxed{0.229}$

e)  $12 \cdot e^{0 \cdot h \cdot t} > 24 \cdot e^{0.229 \cdot t} \Rightarrow e^{0 \cdot h \cdot t} > 2 \cdot e^{0.229 \cdot t} \Rightarrow$

$\ln e^{0 \cdot h \cdot t} > \ln(2 \cdot e^{0.229 \cdot t}) \Rightarrow 0 \cdot h \cdot t > \ln 2 + 0.229 \cdot t \Rightarrow$

$0.171t > \ln 2 \Rightarrow t > \frac{\ln 2}{0.171} \approx \boxed{1053 \text{ hours}}$