

Cálculo Diferencial en exámenes BI – NM (Extracto)

Definición e Interpretación geométrica de la derivada

Mayo 00
P1

Let $f(x) = x^3$.

(a) Evaluate $\frac{f(5+h) - f(5)}{h}$ for $h = 0.1$.

(b) What number does $\frac{f(5+h) - f(5)}{h}$ approach as h approaches zero?

Mayo 02
P1

En la **Figura 1** aparecen las gráficas de las funciones f_1 , f_2 , f_3 , f_4 .

En la **Figura 2** se muestran las gráficas de las derivadas de las funciones de la **Figura 1**, por ejemplo, la derivada de f_1 aparece en el diagrama (d).

Figura 1







Figura 2




Rellene la tabla siguiente efectuando la correspondencia entre cada una de las funciones y su derivada.

Función	Diagrama de la derivada
f_1	(d)
f_2	
f_3	
f_4	

Nov 03
P1La siguiente figura muestra la gráfica de la función $y = f(x)$.Dibuje de forma aproximada, sobre la cuadrícula inferior, la gráfica de $y = f'(x)$.

Nov 04
P1

A continuación se muestra parte de la gráfica de la función periódica f . El dominio de f es $0 \leq x \leq 15$ y el período es 3.




(a) Halle

- (i) $f(2)$;
- (ii) $f'(6,5)$;
- (iii) $f'(14)$.

(b) ¿Cuántas soluciones tiene la ecuación $f(x) = 1$ en el dominio dado?

Mayo 06
P1

The following diagram shows part of the curve of a function f . The points A, B, C, D and E lie on the curve, where B is a minimum point and D is a maximum point.



(a) Complete the following table, noting whether $f'(x)$ is positive, negative or zero at the given points.


	A	B	E
$f'(x)$			

(b) Complete the following table, noting whether $f''(x)$ is positive, negative or zero at the given points.

	A	C	E
$f''(x)$			

May 07
P1

The graph of the function $y = f(x)$, $0 \leq x \leq 4$, is shown below.




(a) Write down the value of

(i) $f'(1)$;

(ii) $f'(3)$.

Nov 08
P1

The diagram shows part of the graph of $y = f'(x)$. The x -intercepts are at points A and C. There is a minimum at B, and a maximum at D.



(a) (i) Write down the value of $f'(x)$ at C.

(ii) **Hence**, show that C corresponds to a minimum on the graph of f , i.e. it has the same x -coordinate.

(b) Which of the points A, B, D corresponds to a maximum on the graph of f ?

(c) Show that B corresponds to a point of inflexion on the graph of f .

Mayo 09
P2

Let $f(x) = x^3 - 4x + 1$.


(a) Expand $(x+h)^3$.

(b) Use the formula $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ to show that the derivative of $f(x)$ is $3x^2 - 4$.

Nov 07
P1

On the axes below, sketch a curve $y = f(x)$ which satisfies the following conditions.

x	$f(x)$	$f'(x)$	$f''(x)$
$-2 \leq x < 0$		negative	positive
0	-1	0	positive
$0 < x < 1$		positive	positive
1	2	positive	0
$1 < x \leq 2$		positive	negative



Recta tangente / Recta Normal

Mayo 00
P1 Find the coordinates of the point on the graph of $y = x^2 - x$ at which the tangent is parallel to the line $y = 5x$.

Mayo 06
P1 Considere la función $f : x \mapsto 3x^2 - 5x + k$.

(a) Escriba $f'(x)$.

La ecuación de la tangente a la gráfica de f en $x = p$ es $y = 7x - 9$. Halle el valor de

(b) p ;

(c) k .

May 07
P1 Sea $f(x) = 3 \cos 2x + \sin^2 x$.

(a) Compruebe que $f'(x) = -5 \sin 2x$.

(b) En el intervalo $\frac{\pi}{4} \leq x \leq \frac{3\pi}{4}$, una normal a la gráfica de f tiene por ecuación $x = k$.
Halle el valor de k .

May 07
P1 Let $f(x) = x^3 - 3x^2 - 24x + 1$.

The tangents to the curve of f at the points P and Q are parallel to the x -axis, where P is to the left of Q.

(a) Calculate the coordinates of P and of Q.

Let N_1 and N_2 be the normals to the curve at P and Q respectively.

(b) Write down the coordinates of the points where

(i) the tangent at P intersects N_2 ;

(ii) the tangent at Q intersects N_1 .

Nov 07
P1 Consider the function $f(x) = 4x^3 + 2x$. Find the equation of the normal to the curve of f at the point where $x = 1$.

Nov 09
P2 Consider the curve with equation $f(x) = px^2 + qx$, where p and q are constants. The point A(1, 3) lies on the curve. The tangent to the curve at A has gradient 8. Find the value of p and of q .

Mayo 10
TZ2
P1#5 Sea $f(x) = kx^4$. El punto P(1, k) pertenece a la curva de f . En P, la recta normal a la curva es paralela a $y = -\frac{1}{8}x$. Halle el valor de k .

Mayo 14
TZ1
P2#7 Let $f(x) = \frac{g(x)}{h(x)}$, where $g(2) = 18$, $h(2) = 6$, $g'(2) = 5$, and $h'(2) = 2$. Find the equation of the normal to the graph of f at $x = 2$.

Cálculo de funciones derivadas

Mayo 03
P1

Sea $f(x) = e^{\frac{x}{3}} + 5 \cos^2 x$. Halle $f'(x)$

Nov 04
P1

Sea $f(x) = 6 \sqrt[3]{x^2}$. Halle $f'(x)$.

Nov 04
P2

Sea $f(x) = 1 + 3 \cos(2x)$ para $0 \leq x \leq \pi$, donde x viene expresado en radianes.

(a) (i) Halle $f'(x)$.

(ii) Halle los valores de x para los cuales $f'(x) = 0$, expresando las respuestas en función de π .

Nov 05
P1

Let $f(x) = (2x + 7)^3$ and $g(x) = \cos^2(4x)$. Find

(a) $f'(x)$;

(b) $g'(x)$.

Mayo 06
P1

(a) Let $f(x) = e^{5x}$. Write down $f'(x)$.

(b) Let $g(x) = \sin 2x$. Write down $g'(x)$.

(c) Let $h(x) = e^{5x} \sin 2x$. Find $h'(x)$.

Mayo 09
P1

Sean $f(x) = e^{-3x}$ y $g(x) = \sin\left(x - \frac{\pi}{3}\right)$.

(a) Escriba

(i) $f'(x)$;

(ii) $g'(x)$.

(b) Sea $h(x) = e^{-3x} \sin\left(x - \frac{\pi}{3}\right)$. Halle el valor exacto de $h'\left(\frac{\pi}{3}\right)$.

Nov 09
P2

Let $f(x) = \cos 2x$ and $g(x) = \ln(3x - 5)$.

(a) Find $f'(x)$.

(b) Find $g'(x)$.

(c) Let $h(x) = f(x) \times g(x)$. Find $h'(x)$.

Muestra
14
P1#7

Given that $f(x) = \frac{1}{x}$, answer the following.

(a) Find the first four derivatives of $f(x)$.


(b) Write an expression for $f^{(n)}(x)$ in terms of x and n .

Mayo 11
TZ2 P1#4

Sea $h(x) = \frac{6x}{\cos x}$. Halle $h'(0)$.

Máximos / Mínimos y Puntos de Inflexión**Mayo 06**
P1


El siguiente diagrama presenta la gráfica de $f(x) = x^2 e^{-x}$ para $0 \leq x \leq 6$. Existen puntos de inflexión en A y C y existe un máximo en B.



- (a) Utilizando la regla del producto para la derivación, halle $f'(x)$.
- (b) Halle el valor **exacto** de la **coordenada y** de B.
- (c) La derivada segunda de f es $f''(x) = (x^2 - 4x + 2)e^{-x}$. Utilice este resultado para hallar el valor **exacto** de la **coordenada x** de C.

Mayo 06
P1

El diagrama siguiente presenta la gráfica de una función g .



La pendiente de la curva tiene un valor máximo en el punto B y un valor mínimo en el punto D. La tangente es horizontal en los puntos C y E.

- (a) Complete la siguiente tabla, indicando si la derivada primera g' es positiva o negativa y si la derivada segunda g'' es positiva o negativa.

Intervalo	g'	g''
$a < x < b$		
$e < x < f$		

- (b) Complete la tabla que aparece a continuación indicando los puntos de la gráfica descritos por las siguientes condiciones.


Condiciones	Punto
$g'(x) = 0, g''(x) < 0$	
$g'(x) < 0, g''(x) = 0$	

Mayo 08
P2

Let $f(x) = e^x(1 - x^2)$.

- (a) Show that $f'(x) = e^x(1 - 2x - x^2)$.

Part of the graph of $y = f(x)$, for $-6 \leq x \leq 2$, is shown below. The x -coordinates of the local minimum and maximum points are r and s respectively.



- (b) Write down the **equation** of the horizontal asymptote.
- (c) Write down the value of r and of s .
- (d) Let L be the normal to the curve of f at $P(0, 1)$. Show that L has equation $x + y = 1$.


Nov 10
P2#7

Sea $f'(x) = -24x^3 + 9x^2 + 3x + 1$.

- (a) En la gráfica de f hay dos puntos de inflexión. Escriba las coordenadas x de estos puntos.
- (b) Sea $g(x) = f''(x)$. Explique por qué en la gráfica de g no hay ningún punto de inflexión.

Mayo 11
TZ1
P2#9

The following diagram shows the graph of $f(x) = e^{-x^2}$.



The points A, B, C, D and E lie on the graph of f . Two of these are points of inflexion.


- (a) Identify the **two** points of inflexion.
- (b) (i) Find $f'(x)$.
- (ii) Show that $f''(x) = (4x^2 - 2)e^{-x^2}$.
- (c) Find the x -coordinate of each point of inflexion.
- (d) Use the second derivative to show that one of these points is a point of inflexion.

Mayo 12

TZ2

P1#10

Sea $f(x) = \frac{x}{-2x^2 + 5x - 2}$, para $-2 \leq x \leq 4$, $x \neq \frac{1}{2}$, $x \neq 2$. A continuación se muestra la gráfica de f .



La gráfica de f presenta un mínimo local en $A(1, 1)$ y un máximo local en B.

- Utilice la regla del cociente para comprobar que $f'(x) = \frac{2x^2 - 2}{(-2x^2 + 5x - 2)^2}$.
- A partir de lo anterior, halle las coordenadas de B.
- Sabiendo que la recta $y = k$ no corta a la gráfica de f , halle los posibles valores de k .

Mayo 13

TZ1

P1#10

Consider $f(x) = \ln(x^4 + 1)$.

- Find the value of $f(0)$.
- Find the set of values of x for which f is increasing.

The second derivative is given by $f''(x) = \frac{4x^2(3 - x^4)}{(x^4 + 1)^2}$.


The equation $f''(x) = 0$ has only three solutions, when $x = 0, \pm\sqrt[4]{3} (\pm 1.316...)$.

- Find $f''(1)$.
 - Hence**, show that there is no point of inflexion on the graph of f at $x = 0$.
- There is a point of inflexion on the graph of f at $x = \sqrt[4]{3} (x = 1.316...)$. Sketch the graph of f , for $x \geq 0$.

Aplicación de la derivada en la resolución de Problemas


Nov 05
P1

The following diagram shows a rectangular area ABCD enclosed on three sides by 60 m of fencing, and on the fourth by a wall AB.




Mayo 08
P1

Find the width of the rectangle that gives its maximum area.
The following diagram shows a semicircle centre O, diameter [AB], with radius 2. Let P be a point on the circumference, with $\angle POB = \theta$ radians.



- Find the area of the triangle OPB, in terms of θ .
- Explain why the area of triangle OPA is the same as the area triangle OPB.

Let S be the total area of the two segments shaded in the diagram below.



- Show that $S = 2(\pi - 2 \sin \theta)$.
- Find the value of θ when S is a local minimum, justifying that it is a minimum.
- Find a value of θ for which S has its greatest value.

Mayo 10
TZ2
P2#7

El número n de bacterias que hay en un plato en el instante t minutos viene dado por $n = 800e^{0.13t}$.

- Halle el valor de n para $t = 0$.
- Halle la tasa de variación a la que está aumentando n en el instante $t = 15$.
- Transcurridos k minutos, la tasa de incremento de n es mayor que 10000 bacterias por minuto. Halle el valor mínimo de k , donde $k \in \mathbb{Z}$.

Mayo 14
TZ1
P2#5


The population of deer in an enclosed game reserve is modelled by the function $P(t) = 210 \sin(0.5t - 2.6) + 990$, where t is in months, and $t = 1$ corresponds to 1 January 2014.

- Find the number of deer in the reserve on 1 May 2014.
- Find the rate of change of the deer population on 1 May 2014.
 - Interpret the answer to part (i) with reference to the deer population size on 1 May 2014.

Cinemática

Mayo 00
P2#2

The main runway at *Concordville* airport is 2 km long. An aeroplane, landing at *Concordville*, touches down at point T, and immediately starts to slow down. The point A is at the southern end of the runway. A marker is located at point P on the runway.



As the aeroplane slows down, its distance, s , from A, is given by

$$s = c + 100t - 4t^2,$$

where t is the time in seconds after touchdown, and c metres is the distance of T from A.

- (a) The aeroplane touches down 800 m from A, (i.e. $c = 800$).
- Find the distance travelled by the aeroplane in the first 5 seconds after touchdown.
 - Write down an expression for the velocity of the aeroplane at time t seconds after touchdown, and hence find the velocity after 5 seconds.

The aeroplane passes the marker at P with a velocity of 36 m s^{-1} . Find

- how many seconds after touchdown it passes the marker;
 - the distance from P to A.
- (b) Show that if the aeroplane touches down before reaching the point P, it can stop before reaching the northern end, B, of the runway.

Nov 00
P2#2

A rock-climber slips off a rock-face and falls vertically. At first he falls freely, but after 2 seconds a safety rope slows him down. The height h metres of the rock-climber after t seconds of the fall is given by:

$$\begin{aligned} h &= 50 - 5t^2, & 0 \leq t \leq 2 \\ h &= 90 - 40t + 5t^2, & 2 \leq t \leq 5 \end{aligned}$$

- Find the height of the rock-climber when $t = 2$.
- Sketch a graph of h against t for $0 \leq t \leq 5$.
- Find $\frac{dh}{dt}$ for:
 - $0 \leq t \leq 2$
 - $2 \leq t \leq 5$
- Find the velocity of the rock-climber when $t = 2$.
- Find the times when the velocity of the rock-climber is zero.
- Find the minimum height of the rock-climber for $0 \leq t \leq 5$.

Mayo 04
TZ2
P1#7

El desplazamiento, s metros, de un coche a los t segundos, respecto a un punto fijo A, viene dado por $s = 4t + 5 - 5e^{-t}$.

- Halle una expresión para la velocidad (en ms^{-1}) a los t segundos.
- Halle la aceleración (en ms^{-2}) en A.

Nov 04
P1#9

El desplazamiento s metros de un coche, t segundos después de pasar por un punto fijo A, viene dado por


$$s = 10t - 0,5t^2.$$

- Calcule la velocidad cuando $t = 0$.
- Calcule el valor de t cuando la velocidad es cero.
- Calcule el desplazamiento del coche respecto a A cuando la velocidad es cero.

Actividades con uso de Calculadora Gráfica

Nov 05
P1

Let $f(x) = 6\sin \pi x$, and $g(x) = 6e^{-x} - 3$, for $0 \leq x \leq 2$. The graph of f is shown on the diagram below. There is a maximum value at B(0.5, b).



- Write down the value of b .
- On the same diagram, sketch the graph of g .
- Solve $f(x) = g(x)$, for $0.5 \leq x \leq 1.5$.

Mayo 06
TZ2
P1#14


El desplazamiento s metros en un tiempo t segundos está dado por

$$s = 5 \cos 3t + t^2 + 10, \text{ para } t \geq 0.$$

- Escriba el valor mínimo de s .
- Halle la aceleración, a , en el tiempo t .
- Halle el valor de t cuando el valor **máximo** de a ocurre por primera vez.

Mayo 08
P2


The following diagram shows the graphs of $f(x) = \ln(3x-2)+1$ and $g(x) = -4\cos(0.5x)+2$, for $1 \leq x \leq 10$.



- Find $f'(x)$.
 - Find $g'(x)$.
- There are two values of x for which the gradient of f is equal to the gradient of g . Find both these values of x .

**Muestra
06 P2**


The diagram below shows the graphs of $f(x) = 1 + e^{2x}$, $g(x) = 10x + 2$, $0 \leq x \leq 1.5$



- (a) (i) Write down an expression for the vertical distance p between the graphs of f and g .
- (ii) Given that p has a maximum value for $0 \leq x \leq 1.5$, find the value of x at which this occurs.

**Mayo 10
TZ2
P2#10**

Considere $f(x) = x \ln(4 - x^2)$, para $-2 < x < 2$. A continuación se muestra la gráfica de f .



- (a) Sean P y Q puntos pertenecientes a la curva de f en los cuales la tangente a la gráfica de f es paralela al eje x .
- (i) Halle la coordenada x de P y de Q.
- (ii) Considere $f(x) = k$. Escriba todos los valores de k para los cuales existen exactamente dos soluciones.

Sea $g(x) = x^3 \ln(4 - x^2)$, para $-2 < x < 2$.

(b) Compruebe que $g'(x) = \frac{-2x^4}{4 - x^2} + 3x^2 \ln(4 - x^2)$.

(c) Dibuje aproximadamente la gráfica de g' .

(d) Considere $g'(x) = w$. Escriba todos los valores de w para los cuales existen exactamente dos soluciones.

**Nov 11
P2#10**

Sea $f(x) = \frac{20x}{e^{0.3x}}$, para $0 \leq x \leq 20$.

- (a) Dibuje aproximadamente la gráfica de f .
- (b) (i) Escriba la coordenada x del punto máximo en la gráfica de f .
- (ii) Escriba el intervalo en el cual f es creciente.
- (c) Compruebe que $f'(x) = \frac{20 - 6x}{e^{0.3x}}$.
- (d) Halle el intervalo en el cual la razón de cambio de f es creciente.